

March 3, 2010

A Multi-wavelength analysis of M81 (NGC3031): insight on the nature of Arp's loop

A. Sollima¹, A. Gil de Paz², D. Martinez-Delgado^{1,3}, R. J. Gabany⁴, J. Gallego⁵, and T. Hallas⁶

¹Instituto de Astrofisica de Canarias, La Laguna, E38205, Spain

²Departamento de Astrofisica, Universidad Complutense de Madrid, E28040, Spain

³Max-Planck-Institut fur Astronomie, Heidelberg, Germany

⁴Blackbird Observatory, New Mexico, USA

⁵Fosca Nit Observatory, Monsec Astronomical Park Observatory, Ager, Spain

⁶Hallas Observatory Annex, Foresthill, USA

DRAFT

ABSTRACT

Context: The optical ring like structure detected by Arp (1965) around M81 (commonly referenced as "Arp's loop") represents one of the most spectacular features observed in nearby galaxies. Arp's loop is commonly interpreted as a tail resulting from the tidal interaction between M81 (NGC3031) and M82 (NGC3034). However, since its discovery the nature of this feature has remained controversial.

Aims: Our primary purpose was to identify the sources of optical and infrared emission observed in Arp's loop.

Methods: The morphology of the Arp's loop has been investigated with deep wide-field optical images. We also measured its colors using IRAS and Spitzer-MIPS infrared images then

compared them with images of M81 (NGC3031) and galactic dust cirrus that fills the area of the sky where M81 (NGC3031) is located.

Results: Optical images reveal that this peculiar object has a filamentary structure characterized by many dust features overlapping M81's field of view. The ratios of far-infrared flux and the estimated dust-to-gas ratios indicate the infrared emission of Arp's loop is dominated cold dust that is most likely from Galactic cirrus that lies above the Milky Way.

Conclusions: The above results suggest the light observed at optical wavelengths is a combination of emission from i) a few recent star forming regions located close to M81 (NGC3031) where both bright UV complexes and peaks in the HI distribution are found, ii) the extended disk of M81 and iii) scattered light from the same Galactic cirrus that is responsible for the bulk of the far infrared emission.

Key words. Methods: data analysis – Techniques: photometric – galaxies: individual: M81 – infrared: galaxies

1. Introduction

The nearby galaxy M81 (NGC3031) together with the galaxies M82 (NGC3034), NGC3077 and NGC2976 forms one of the most iconic groups of interacting galaxies. Located at a distance of ~ 3.7 Mpc (Makarova et al. 2002), the M81 group has been a subject for many studies analyzing evidence of strong interactions among its components. The group contains remnants of tidal bridges connecting the three most prominent galaxies, visible in HI survey (Gottesman & Weliachew 1975; Yun et al. 1994), and over 40 dwarf galaxies (Karachentsev et al. 2001). In 1965, Halton Arp detected an unusual ring shaped feature in the vicinity of M81 (NGC3031) while examining photographic plates produce with the 48-inch Schmidt on Mount Palomar. This

optical feature (usually referenced as "Arp's loop") is located 17 arcmin north-east of M81's center and covers a wide area of ~ 160 square arcmin. It also exhibits symmetry towards the end of the M81 disk, slightly tilted toward M82.

Typically, Arp's loop is interpreted as a tail resulting from the tidal interaction between M81 (NGC3031) and M82 (NGC3034). However, since its discovery this interpretation has been doubted by many authors. In fact, the region containing the M81 group of galaxies is filled by Galactic cirrus that covers a large area of the sky near M81 (Sandage 1976). Arp's loop exhibits color and emission properties similar to those observed in Galactic cirrus clouds (Abolins & Rice 1987; Appleton et al. 1993; Bremnes et al. 1998) and this has raised doubts about its link with M81.

Following its discovery, studies supporting the extra-Galactic nature of Arp's loop were based on 21cm HI observations in the M81 group region (Yun et al. 1994). These authors detected a collimated emission in the direction of the north-eastern part of the Arp's loop which blends smoothly into the structure and velocity of the HI disk of M81 (NGC 3031). Numerical simulations of the system by Yun (1999) successfully reproduced the HI tidal debris assuming M82 (NGC3034) and NGC 3077 approached M81 during recent epochs.

More recently, Makarova et al. (2002) and De Mello et al. (2008) resolved the nebular region, characterized by strong HI emission, into stars using deep HST observations. Their color-magnitude diagrams (CMD) showed the presence of a significant young star population (with an age of $\sim 40 - 160$ Myr) and an old population characterized by a well defined Red Giant Branch. The magnitude of the tip of the Red Giant stars assumed to be associated with Arp's loop indicated a distance comparable with that of M81 (NGC3031) (Karachentsev et al. 2002). These authors suggested that the old component could have been removed from the M81 and/or M82 disk during their mutual interaction.

Further insight was recently provided by Barker et al. (2009). This team analyzed a wide area that included M81 (NGC3031) using deep Suprime-Cam observations. Their results revealed no overdensity of Red Giant stars along Arp loop's extension (which is at odds with De Mello et al.'s results), whereas a significant group of young Red Supergiants and Main Sequence stars were evident in the confined region previously surveyed in HST studies. These authors concluded that Arp's loop was originally a gaseous tidal debris stream that formed stars only in the last 200-300 Myr. The same qualitative results have also been obtained by Davidge (2009).

In this paper we present the results of a multi-wavelength analysis of Arp's loop using the deepest optical and infrared observations available.

2. The Data

The analysis presented here is based on two datasets: i) a set of deep ground based optical images obtained at different facilities (the Fosca Nit Observatory and Hallas Observatory) and ii) a set of far-infrared images observed with MIPS at the Spitzer Space Telescope at $24\mu\text{m}$, $70\mu\text{m}$ and $160\mu\text{m}$ and with IRAS using the latest IRIS data products.

The first dataset was collected with a commercially available 106mm aperture f/5 Takahashi FSQ refractor telescope of the Fosca Nit Observatory (FNO) situated near Ager (Spain) at the Monsec Astronomical Park. We used a Santa Barbara Imaging Group (SBIG) STL-11000M CCD camera which yields a large field of view (3.9×2.7) at a plate scale of 3.5 arcsec/pixel. The image set consists of multiple deep exposures through four Optec Inc. broadband (LBGR) filters. A set of individual 600 sec images were obtained during several photometric nights between January and February 2008, achieving a total exposure times of the co-added images of 17,100 sec.

High resolution images were also gathered, during four photometric nights between February and March 2007, with a 14.5" f/8 RCOS cassegrain telescope situated at the Hallas Observatory Annex (HOA) situated near Foresthill, California (USA). A Santa Barbara Imaging Group STL-11000M camera, yielding a $41' \times 27'$ field of view and a 1 arcsec/pixel plate scale, was attached to the Cassegrain focus of the telescope. A set of images through four broadband (LBGR) filters have been secured and co-added, yielding a total exposure time of 51,400 sec.

The images were reduced following standard procedures for bias correction and flat-fielding. To enhance the signal-to-noise of the faint structures around M81, image noise was attenuated with a Gaussian filter (Davis 1990).

MIPS images were also retrieved from the Spitzer Infrared Nearby Galaxy Survey (SINGS Data Release 4, Kennicutt et al. 2003) public archive. The sample consists of a set of deep images in the 24 μ m, 70 μ m and 160 μ m bands. The total exposure time is approximately 220 s, 84 s and 25 s at 24 μ m, 70 μ m and 160 μ m, respectively. Individual frames have been reduced using the MIPS Instrument Team Data Analysis Tool (Gordon et al. 2005). The background was subtracted using the average value of an empty region of the MIPS field of view. The uncertainties of the final absolute calibrations were estimated at 10% for the 24 μ m data and 20% for both the 70 μ m and 160 μ m data. The combined images were then aligned in the standard WCS reference frame. The overall field of view is approximately 30'x44'.

To compare the colors of Arp's loop with those of nearby Galactic cirrus we also retrieved the newest generation IRIS imaging products from the IRAS satellite covering a large region (12.5.x 12.5.) around M81 (see Figure 3). The IRIS images that we used benefit from better zodiacal-light subtraction, calibration, zero levels, and destripping than previous versions. In particular, the 100 μ m IRIS maps represent a significant improvement over those used by Schlegel et al. (1998).




Fig. 1. Upper panel: 2. \times 3. image of the M81 region from Witt et al. (2008; by courtesy of S. Mandel). North is up, east to the right. Bottom panel: FNO image of the M81 group of galaxies.

3. Results

3.1. Optical imaging

Figure 1 (lower panel) shows a wide field view around M81 (NGC3031) obtained with the FNO telescope. This image clearly shows many large- scale cirrus structures. The width of these filaments range from ~ 30 arcsec up to several arcmin and extend in connective patterns over several degrees (as it is clearly visible in the panorama of this sky region displayed in the image obtained by Witt et al. 2008; see Fig.1, top panel).

Figure 2 displays the image obtained by combining all the HOA observations. This image covers an area of $41' \times 27'$. This deep image reaches a point-source limiting visual magnitude of $V \sim 27$. The spectacular structure of Arp's loop is evident in the north-eastern region. Arp's loop appears as a dim filamentary ring that wraps and overlaps the galaxy's disk. A careful inspection of M81's disk reveals the presence of several dust absorption features that correspond with the intersection of Arp's loop and the galaxy. These features, already noticed by Arp (1965) and Efremov et al. (1986), are due to the presence of dust in the ring, indicating that part of it should be situated between the observer and M81. In the north-eastern part of this feature (in the region where HI strong emission was found by Yun et al. 1994) a significant overdensity of stellar complexes is also apparent and suggests that new stars are still forming (Makarova et al. 2002; Mouhcine & Ibatá 2010).




Fig. 2. Combined HOA image of the M81 region. The field of view of this image is $41' \times 27'$. The North and east directions are indicated. Arp's loop is visible as the nebular ring crossing the disk of M81 on the right side of the image. Dust absorption features are superimposed over and surround the disk of M81.

3.2. Far Infrared emission

As previously mentioned in Sect. 1, far infrared imaging is a fundamental tool to highlight emission from dusty objects. This is particularly important when analyzing Arp's loop since both dust absorption in the form of dust lanes projected against the disk of M81 and dust emission from the loop itself, are clearly seen (see Sect. 3.1). The right panel of Figure 3 shows an IRAS map in a wide region (12.5×12.5) around the M81 group of galaxies constructed using images in the $12\mu\text{m}$, $60\mu\text{m}$ and $100\mu\text{m}$ bands. As noted in Figure 1, the surveyed area is filled with Galactic cirrus clouds.

To compare Arp's loop with the cirrus clouds in infrared light, we measured the flux for various regions of the $60\mu\text{m}$ and $100\mu\text{m}$ IRAS image (see Figure 3). Removal of the background (mainly of cosmic origin, since zodiacal light had been previously removed from the IRIS images) was performed using the minimum value of the surface brightness in the 12.5×12.5 . IRIS image for each band. Despite its admittedly large amount of uncertainty, this method has been already

used in the past (Carey et al. 1997) and yields a value for the background at 100 μm (0.67 MJy/sr) which is very similar to the average Cosmic Infrared Background of 0.78 MJy/sr as determined by Lagache et al. (2000). The selected regions, the adopted apertures and the resulting flux for M81 (NGC3031) and Arp loop's regions are listed in Table 1.

In the left panel of Figure 3, the ratio between the 60 μm and 100 μm flux are plotted for the main body of M81 (open black point), cirrus clouds (grey points) and Arp's loop (black filled points) as a function of the 100 μm surface brightness. It should be noted that cirrus clouds occupy a well defined sequence in this diagram, with high-density cirrus showing a lower ratio than the most diffuse cirrus in the field. This trend has been previously reported by several authors (e.g. Abergel et al. 1994). For comparison, the "reference value" commonly adopted for the I60 μm /I100 μm ratio in high latitude cirrus is ~ 0.2 (see e.g. Arendt et al. 1998). M81 presents a high color (I60 μm /I100 μm ~ 0.3) compared to Galactic cirrus of the same brightness, although it is not very different from neither diffuse cirrus colors nor from that of the loop itself. In this sense, it is worth noting that M81 (NGC3031) is a very quiescent object in terms of its dust emission properties (see e.g. Dale et al. 2007). In fact, it has the lowest I60 μm /I100 μm ratio among those in the Helou (1986) sample of galaxies. More importantly, the colors found within the circular apertures placed on the loop (cyan circles in Figure 3) are not very different from the color of the M81 disk and follow the same trend as the cirrus. This latter fact suggests a Galactic origin for the bulk of the far infrared emission associated with Arp's loop.

Despite these results, it is still conceivable that dust in the outer disk of M81 (NGC3031), or in a potential tidal feature around M81, could show the same colors (and follow the same surface brightness trend) as the cirrus because they could contain cold dust with emission properties similar to Galactic cirrus. The reader is referred to the recent work by Bot et al. (2009) for an extensive discussion on the properties of Galactic cirrus.

A deeper insight into the structure of dust emission around M81 (NGC3031) necessarily requires the use of deep high-resolution (arcmin-scale) Spitzer-MIPS observations. The MIPS 160 μm data contains information at longer wavelengths that might offer further clues about the possible differences (or similarities) between dust emission in the loop and M81 (NGC3031). The right panel of Figure 4 displays a color emission map from Spitzer. To construct this map, the MIPS/Spitzer 24 μm and 70 μm images were binned and scaled to the resolution of the 160 μm image using convolution kernels developed by Gordon et al. (2008). To compare the far infrared colors of Arp's loop with the stellar component of M81 (NGC3031), we measured

the 70 μm and 160 μm flux sampled in circular apertures positioned at four regions of Arp’s loop and across the M81’s disk (see the right panel of Fig. 4). We exercised care in selecting both regions populated by stellar complexes and regions where only an extended diffuse emission is noticeable from optical images along Arp’s loop. The measured flux are listed in Table 1. Most importantly, the 70 μm /160 μm flux ratio as a function of the 160 μm brightness is shown in the left panel. Note that Arp’s loop and the disk of M81 (NGC3031) displays very different colors. Here the segregation between M81 (NGC3031) and the loop is much clearer than it is in the IRAS I60 μm /I100 μm ratio. In particular, the ratio 70 μm /160 μm appears significantly smaller (<0.1 in all cases) than that measured on the galaxy’s disk (which typically ranges between 0.2-0.4 in the inner disk and form a plateau around 0.12 in the outer disk) along the entire extension of the loop. The 70 μm /160 μm ratios measured in the loop are also similar to the 60 μm /160 μm ratios found by Bot et al. (2009) in their analysis of serendipitous observations of Galactic cirrus contained in the SINGS Spitzer-MIPS fields.




Fig. 3. In the left panel, the I60 μm /I100 μm ratio is shown as a function of the I100 μm brightness for the loop (black filled points), M81 (open point) and the surrounding cirrus features (grey points). The typical uncertainty of the regions along Arp’s loop is shown in the bottom part of the panel. The right panel depicts a false-color IRAS image of the M81(NGC3031) region. North is up, east is to the left. The circular apertures used in the analysis of the Galactic cirrus (yellow circles; 12’ diameter), M81 (green circle; 12’ diameter) and Arp’s loop (tiny cyan circles above M81; 4’ diameter) are shown.




Fig. 4. In the left panel, the $I_{70\mu\text{m}}/I_{160\mu\text{m}}$ ratio calculated for Arp's loop (filled circles) and the M81 (NGC3031) disk (open circles) are shown as a function of the $I_{160\mu\text{m}}$ brightness. The typical uncertainty is shown in the upper part of the panel. In the right panel the $27' \times 41'$ color map of the Spitzer MIPS emission for M81 was constructed using the $24\mu\text{m}$, $70\mu\text{m}$ and $160\mu\text{m}$ images. The circular apertures are also shown with the same color code used in Figure 3.

3.3. Comparison between Far-Infrared and HI emission

Figure 5 displays a map of the HI emission measured by Walter et al. (2008) and the MIPS $160\mu\text{m}$ flux over plotted on the optical HOA image. Notice that the HI emission nicely follows the spiral arms of M81 (NGC3031). Also, Holmberg IX (visible on the east of M81 in Figure 5) and the southern part of Arp's loop are embedded in clouds connected to the main body of M81 (NGC3031). On the other hand, most of the far infrared emission appears to be associated to the disk of M81 (NGC3031), with a significant contribution from Arp's loop. In addition, note that the HI emission extends across the entire galaxy disk well beyond its optical extent, partially overlapping Arp's loop. It is worth noting that while the $160\mu\text{m}$ flux is relatively high along the entire loop extension, the HI emission disappears in the northern part. To better visualize this comparison, the contours of the HI emission measured by Walter et al. (2008) are over plotted to the MIPS $160\mu\text{m}$ image in Fig. 6. The mismatch between the spatial distribution of the HI and MIPS $160\mu\text{m}$ emission suggests that different regions of Arp's loop are characterized by different emitting mechanisms.

A more quantitative comparison has been performed by estimating the relative fraction of dust and gas over Arp loop's extension and across the main body of M81 (NGC3031). This parameter is particularly powerful in discriminating between the "canonical" emission of galactic disks and other dust-dominated sources. Indeed, this quantity spans a limited range among the observed galaxies and it has been found to show a well defined radial behavior, decreasing at large distances from the galactic center (Issa et al. 1990; Boissier et al. 2004; Munoz-Mateos et al. 2009).

To estimate the total dust mass, we used MIPS flux at 24 μ m, 70 μ m and 160 μ m measured in the same circular apertures along Arp's loop and across the M81's disk defined in Sect. 3.2 (see also the right panel of Fig. 7). We adopted the relation by Munoz-Mateos et al. (2009; see their eq. A8) to convert flux in mass adopting a distance for M81 of 3.7 Mpc (Makarova et al. 2002). In order to obtain total gas mass, we measured the HI flux from the M81 (NGC3031) intensity map obtained by the THINGS survey (Walter et al. 2008). For this purpose the THINGS image has been binned and scaled to the resolution of the 160 μ m image using the convolution kernels developed by Gordon et al. (2008). The corresponding total gas mass has been calculated using eq. 3 of Walter et al. (2008). The resulting dust and HI mass together with the corresponding dust-to-gas ratios (DGR) are listed in Table 1.

Equivalent radial distances from the galactic center have been measured by assuming the position angle and axial ratio defined by Munoz-Mateos et al. (2009). In the left panel of Figure 7, the obtained DGR values are plotted against the equivalent angular radius along the semi-major axis. The observed profile of M81 (NGC3031) calculated by Munoz-Mateos et al. (2009) is over plotted for comparison. It is evident that at distances $r' > 200''$ the DGR shows the typical decreasing trend with galactocentric distance. Note that the DGR values measured along Arp's loop are significantly larger than that measured in the central region of M81 (NGC3031) and deviate from the general radial trend.

It is interesting to note that the maximum DGR ($\log(M_{\text{dust}}/M_{\text{gas}}) \sim 0.12$) is found in the northern side of Arp's loop, where redder MIPS colors have been measured and the HI emission drops below the detection limit. Such a large ratio has never been observed in any galaxy analyzed by Munoz-Mateos et al. (2009) regardless of the distance from the galactic center.

On the other hand the regions along Arp's loop with smaller DGR values are those located in the north-eastern part of the loop (where UV complexes have been observed; De Mello et al. 2008) and in the northwestern part (where the spiral arm of M81(NGC3031) overlaps Arp's loop). The surprisingly high DGR derived along Arp's loop results from the significantly small HI flux measured on the Walter et al. (2008) maps.

To check the dependence of the obtained results on the adopted dataset, we also calculated the DGR using the HI map provided by Yun et al. (1994). As apparent in Fig. 7, the derived DGR are very similar to those obtained using the Walter et al.'s map. Smaller intensities of the 21cm emission in the eastern and northern part of Arp's loop are also evident in the single-dish map obtained by Appleton & van der Hulst (1988). Moreover, the HI map of Walter et al. (2008) has been constructed using VLA data decuded in both C and D configurations. On the basis of all these considerations, we concluded that the large measured DGR can be assigned to sensitivity problems and/or filtering by the interferometer of the Walter et al. (2008) data.




Fig. 5. HOA images of the region around M81 (NGC3031). The HI emission from Walter et al. (2008) (left panel) and the MIPS 160.03bcm image (right panel) has been overlotted.




Fig. 6. MIPS 160 μ m image of M81 (NGC3031). The contours of the HI emission from Walter et al. (2008) have been over plotted.

The large DGR measured along Arp's loop supports the idea that the far-infrared emission is increased due to the contribution of a dust-dominated source. In this regard, the hypothesis about the presence of Galactic cirrus fit this picture since it would provide a significant contribution to the far-infrared light filtering and HI emission.

4. Discussion

This paper presents a multi-wavelength analysis of Arp's loop using the deepest optical and infrared images available.

Arp's loop appears to be formed by a filamentary wrap that partially overlaps the disk of M81 (NGC3031). The many dust absorption features overlapping M81's field of view seen in the HOA images suggests that part of it is situated between the observer and the galaxy.

The comparison between the ratio of far infrared flux emitted at 70 μ m and 160 μ m by Arp's loop and M81 (NGC3031) indicates that the emission from the loop is dominated by the contribution of cold dust. Its far infrared emission resembles the properties of Galactic cirrus clouds, which also share the same colors in the IRAS bands. The same conclusion is also supported by the analysis of the DGR measured along Arp loop's extension which indicates a surprisingly large relative fraction of dust if compared with that expected at such a large distance from the galactic center. It is therefore likely that at least part of the ring like structure that forms Arp's loop is constituted by Galactic cirrus overlaying the disk of M81 (NGC3031). The region around M81 (NGC3031) is indeed known to be an area of great confusion between Galactic and extra-Galactic objects. Of course, our observations cannot exclude the presence of stripped material from the disk of M81 (NGC3031) and/or M82 (NGC3034) in those regions of Arp's loop where small 170 μ m/160 μ m ratios and high DGR have been measured. However, in this case a selective process of preferentially stripping dust should be performed. Also in this scenario, contamination from the surrounding Galactic cirrus would help to explain all the observational evidence if it were overlapping Arp's loop.

Table 1. Far-Infrared Flux and dust-to-gas ratios

location	RA h m s	Dec deg m s	diameter '	$F_{60\mu m}^{IRIS}$ Jy	$F_{100\mu m}^{IRIS}$ Jy	$F_{70\mu m}^{MIPS}$ Jy	$F_{160\mu m}^{MIPS}$ Jy	M_{dust} $10^7 M_{\odot}$	M_{gas} $10^7 M_{\odot}$	$\log(M_{dust}/M_{gas})$
Arp's loop NW	09 54 34	+69 16 54	4	0.962	3.493	0.285	4.386	0.393	9.043	-1.362
Arp's loop N	09 55 43	+69 19 28	4	0.953	2.953	0.037	2.171	2.075	1.586	0.117
Arp's loop NE	09 57 02	+69 18 56	4	0.903	2.877	0.106	3.670	1.390	14.207	-1.009
Arp's loop E	09 57 09	+69 11 21	4	0.853	2.788	0.198	3.048	0.271	2.6476	-0.990
M81's disk	09 55 33	+69 03 56	12	20.101	65.550	71.98	272.300	2.520	175.790	-2.638




Fig. 7. In the left panel, the DGR calculated for Arp's loop and the M81 (NGC3031) disk are shown as a function of the equivalent angular distance from the galactic center. HI mass has been calculated using both Walter et al. (2008; black points) and Yun et al. (1994; grey points) HI maps. The radial DGR profile measured by Munoz-Mateos et al. (2009) is over plotted. In the right panel, the THINGS image of M81 (NGC3031) is shown. The circular apertures are also shown with the same color code used in Figure 3.

An unambiguous distinction between these two possibilities is very complex (see e.g. Cortese et al. 2010) and cannot be made solely on the basis of morphological arguments or in terms of optical colors. In fact, cirrus clouds and galaxies have largely overlapping colors (Bremnes et al. 1998) and emissions that encompass a wide range of wavelengths from the UV to the far infrared (Haikala et al. 1995).

In this context, it is important to distinguish between the stellar populations surrounding M81 (NGC3031) and the nebular region which constitutes Arp's loop. In recent years, many sites of recent star formation have been reported in M81's outer disk by GALEX (de Mello et al. 2008). Given the wide area covered by Arp's loop, it is possible that some of these star formation regions could be accidentally located along the direct line of sight path to this feature. In this

respect, the spatial correlation between the HI emission and Arp's loop provides some interesting evidence. As previously mentioned in Sect. 1, these HI clouds share similar dynamics with the disk of M81 (NGC3031). The deep photometric analyses of this region performed by Karachentsev et al. (2002), Makarova et al. (2002), Sun et al. (2005), De Mello et al. (2008), Davidge (2009) and Barker et al. (2009) revealed evidence for a young stellar population located at a distance comparable to M81 (NGC3031). Moreover, De Mello et al. (2008) identified eight FUV sources in this region using GALEX images. The presence of a stellar component associated with M81 (NGC3031) in this area, therefore, appears indisputable. However, notice that the HI emission disappears in the northern part of the loop where both optical and infrared images show significant contributions. Moreover, the analysis of CMDs obtained by Davidge (2009) and Barker et al. (2009) has not detected any overdensity of Red Giants along Arp's loop and only identified a population of young stars in the confined region where the strong HI emission was observed. Thus, although the origin of part of the optical and UV emission in Arp's loop could be emitted by recent star formation episodes, most likely part of it has a Galactic origin.

It is also noticeable that both HI and GALEX emissions extend across the entire disk of M81(NGC3031) well beyond its optical cutoff, partially overlapping the north-western part of Arp's loop (see also Gil de Paz et al. 2007; Thilker et al. 2007). This was also confirmed by the analysis of HST CMDs which revealed the presence of the M81's stellar disk population where GALEX and far infrared emissions overlap (Gogarten et al. 2009).

The observational evidence put forward in this paper suggests that the structure known as Arp's loop is likely formed by three distinct components: i) one is associated with the M81 system that causes the emission detected in the UV and HI plus part of emission seen at optical wavelengths, thus dominating the morphology of the north-eastern part of the loop, ii) a second is dominated by M81's extended disk which contributes to the UV and HI emission in the north-western section of Arp's loop and, iii) another of Galactic origin that dominates the far infrared emission and is responsible for the optical morphology (through scattering by dust) of the entire ring-like structure.

Acknowledgements

This research was supported by the Instituto de Astrofísica de Canarias. We thank the anonymous referee for his/her useful comments and suggestions. We are grateful to Ignacio Trujillo, Michele Bellazzini, Julianne Dalcanton, and Juan Carlos Muñoz-Mateos for helpful

discussions. We also thank Steve Mandel for providing us their wide field deep images of the M81 sky area. We also acknowledge funding from the Spanish Ministry of Education and Science (research project 2007-65090).

References

- Abergel, A., Boulanger, F., Mizuno, A., & Fukui, Y., 1994, *ApJ*, 423, L59
Abolins, J. A., & Rice, W. L., 1987, NASA Conference Publication, 2466, 107
Appleton, P. N., Siqueira, P. R., & Basart, J. P., 1993, *AJ*, 106, 1664
Appleton, P. N., & van der Hulst, J. M., 1988, *MNRAS*, 234, 957
Arendt, R. G., et al., 1998, *ApJ*, 508, 74
Arp, H., 1965, *Science*, 148, 363
Barker, M., Ferguson, A. M., Irwin, M., Arimoto, N., & Jablonka, P., 2009, *AJ*, 138, 1469
Boissier, S., Boselli, A., Buat, V., Donas, J., & Milliard, B., 2004, *A&A*, 424, 465
Bot, C., et al., 2009, *ApJ*, 695, 469
Bremnes, T., Binggeli, B., & Prugniel, P., 1998, *A&AS*, 129, 313
Carey, S. J., Shipman, R. F., Clark, F. O., & Assendorp, R., 1997, *ApJ*, 479, 303
Cortese, L., Bendo, G. J., Isaak, K. G., Davies, J. I., & Kent, B. R., *MNRAS*, in press, ArXiv 1001.098
Dale, D. A., et al., 2007, *ApJ*, 655, 863
Davidge, T. J., 2009, *ApJ*, 697, 1439
Davies, E. 1990, in *Machine Vision: Theory, Algorithms and Practicalities*, (Academic Press), p. 42
de Mello, D. F., et al., 2008, *AJ*, 135, 548
Efremov, Y. N., Karachentsev, I. D., & Karachentseva, V. E., 1986, *Pis ma Astronomicheskii Zhurnal*, 12, 434
- Gil de Paz, A., et al., 2007, *ApJS*, 173, 185
Gogarten, S. M., et al., 2009, *ApJ*, 691, 115
Gordon, K. D., et al., 2005, *PASP*, 117, 503
Gordon, K. D., et al., 2008, *ApJ*, 682, 336
Gottesman, S. T., & Weliachew, L., 1975, *ApJ*, 195, 23
Haikala, L. K., et al., 1995, *ApJ*, 443, L33
Helou, G., 1986, *ApJ*, 311, L33
Issa, M. R., MacLaren, I., & Wolfendale, A. W., 1990, *A&A*, 236, 237
Karachentsev, I. D., Karachentseva, V. E., & Huchtmeier, W. K., 2001, *A&A*, 366, 428
Karachentsev, I. D., et al., 2002, *A&A*, 383, 125

Kennicutt, R. C., Jr., et al. 2003, PASP, 115, 928
Lagache, G., Haffner, L. M., Reynolds, R. J., & Tufte, S. L., 2000, A&A, 354,247
Makarova, L. N., et al. 2002, A&A, 396, 473
Mouhcine, M., & Ibata, R., 2010, MNRAS, in press (arXiv:1002.0456)
Muñoz-Mateos, J. C., et al., 2009, ApJ, 701, 1965
Sandage, A., 1976, AJ, 81, 954
Schlegel, D. J., Finkbeiner, D. P., & Davis, M., 1998, ApJ, 500, 525
Sun, W.-H., et al., 2005, ApJ, 630, L133
Thilker, D. A., et al., 2007, ApJS, 173, 538
Walter, F., et al., 2008, AJ, 136, 2563
Witt, A. N., Mandel, S., Sell, P. H., Dixon, T., & Vihj, U. P., 2008, ApJ, 679, 497
Yun, M. S., 1999, in Proceedings of IAU Symposium, 186, Galaxy Interactions at Low and High Redshift, ed. J. E. Barnes, and D. B. Sanders, 81

Yun, M. S., Ho, P. T. P., & Lo, K. Y., 1994, Nature, 372, 530